

2016 Recommended Immunizations for Children from Birth Through 6 Years Old



Age	RV	DTaP	Hib	PCV	IPV
Birth	HepB				
1 month	HepB				
2 months	RV	DTaP	Hib	PCV	IPV
4 months	RV	DTaP	Hib	PCV	IPV
6 months	RV	DTaP	Hib	PCV	IPV
12 months	HepB				
15 months	HepB				
18 months	HepB				
19-23 months	HepB				
2-3 years	HepB				
4-6 years	HepB				

Is your family growing? To protect your new baby and yourself against whooping cough, get a Tdap vaccine in the third trimester of each pregnancy. Talk to your doctor for more details.

Shaded boxes indicate the vaccine can be given during shown age range.

SEE BACK PAGE FOR MORE INFORMATION ON VACCINE-PREVENTABLE DISEASES AND THE VACCINES THAT PREVENT THEM.

NOTE: If your child misses a shot, you don't need to start over, just go back to your child's doctor for the next shot. Talk with your child's doctor if you have questions about vaccines.

FOOTNOTES: * Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting an influenza (flu) vaccine for the first time and for some other children in this age group.
 † Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 to 18 months later. HepA vaccination may be given to any child 12 months and older to protect against HepA. Children and adolescents who did not receive the HepA vaccine and are at high-risk, should be vaccinated against HepA.

If your child has any medical conditions that put him at risk for infection or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need.